



PHYSICAL EXAM

HEALTH SERVICES 401-232-6220

STUDENT'S NAME: _____ DATE OF BIRTH: _____
 STUDENT ID #: _____
 Date of Exam: _____ (MUST be within one year of university entry - - six months for athletes)

❖ **VITALS MUST BE COMPLETED**

Blood Pressure: _____ Pulse: _____ Height (inches): _____ Weight (lbs): _____ BMI: _____

NORMAL	Check each item in appropriate column Enter NE if not evaluated	NOTE: Describe every abnormality in detail
	Head, face, neck, scalp	
	Nose and sinuses	
	Mouth, teeth, throat	
	Ears	
	Eyes	
	Ophthalmoscopic	
	Neck, thyroid	
	Thorax and breasts	
	Lungs	
	Heart	
	Abdomen	
	Anus and rectum	
	Endocrine system	
	G.U. system	
	Upper extremities	
	Lower extremities	
	Feet	
	Spine	
	Neurologic	
	Psychiatric evaluation	
	Skin	
	Lymphatic system	
	Vascular system	
	Other:	

Is this student receiving or does he/she require continuing medical care, therapy, or observation? _____
 If **YES**, please explain (include notation of medication and dosages and plans concerning illness such as heart disease, asthma, diabetes, seizure disorders, etc.) _____

Does this student have any life-threatening allergies? YES ____ NO ____
 _____ (If yes, please send allergic response and suggested treatment)

I hereby declare that the above-named student is medically cleared to participate in intercollegiate athletics at Bryant University.
 Fit to participate YES ____ NO ____

HEALTH CARE PROVIDER:

Print Name: _____ Phone: _____
 Address: _____
 Signature: _____ Date: _____

IMMUNIZATION RECORD

HEALTH SERVICES
401-232-6220

* To be completed and signed by health care provider *

STUDENT'S NAME: _____ DOB: _____
STUDENT ID #: _____

REQUIRED IMMUNIZATIONS (prior to matriculation)

All information must be in English (dates must include month, day, and year)

A. TETANUS, DIPHTHERIA, PERTUSSIS (MUST be within 10 years)

TDAP ____/____/____

B. MMR (MEASLES, MUMPS, RUBELLA) - 2 doses required

Dose 1 given at age 12 months or later - - Dose 2 given at least 28 days after first dose

Dose #1 ____/____/____ Dose #2 ____/____/____

OR

MEASLES (Rubeola) positive immune titer date ____/____/____

RUBELLA (German measles) positive immune titer date ____/____/____

MUMPS positive immune titer date ____/____/____

C. HEPATITIS B - 3 doses required

Dose 2 must be given one month after dose 1. Dose three must be given five months after dose 2.

Dose #1 ____/____/____ Dose #2 ____/____/____ Dose #3 ____/____/____

OR

Hepatitis B titer Reactive _____ non-reactive _____ Date ____/____/____

D. VARICELLA (Chickenpox) - 2 doses required

1. History of chickenpox disease Yes _____ Date ____/____/____

OR

1. Dose #1 ____/____/____ Dose #2 ____/____/____

OR

2. Varicella titer Date ____/____/____ Reactive _____ non-reactive _____

E. MENINGOCOCCAL QUADRIVALENT (Meningitis) - 2 doses, if dose 1 given before age 16

Dose #1 ____/____/____ Dose #2 ____/____/____

F. COVID-19 (Please upload a copy of your Covid-19 vaccination card)

Dose #1 ____/____/____ Pfizer _____ Moderna _____ J&J _____ Other _____

Dose #2 ____/____/____ Pfizer _____ Moderna _____ J&J _____ Other _____

BOOSTER (Not Required)

Dose #3 ____/____/____ Pfizer _____ Moderna _____ Other _____

HEALTH CARE PROVIDER:

Print Name: _____ Phone: _____

Signature: _____ Date: _____

Part I: Tuberculosis (TB) Screening Questionnaire (to be completed by incoming students)

Please answer the following questions:

Have you ever had close contact with persons known or suspected to have active TB disease? Yes No

Were you born in one of the countries or territories listed below that have a high incidence of active TB disease? (If yes, please CIRCLE the country, below) Yes No

Afghanistan	Congo	Iran (Islamic Republic of)	Namibia	Singapore
Algeria	Côte d'Ivoire	Iraq	Nauru	Solomon Islands
Angola	Democratic People's Republic of	Kazakhstan	Nepal	Somalia South Africa
Anguilla	Korea	Kenya	Nicaragua	South Sudan
Argentina	Democratic Republic of the	Kiribati	Niger	Sri Lanka
Armenia	Congo	Kuwait	Nigeria	Sudan
Azerbaijan	Djibouti	Kyrgyzstan	Northern Mariana	Suriname
Bangladesh	Dominican Republic	Lao People's Democratic	Islands	Swaziland
Belarus	Ecuador	Republic	Pakistan	Tajikistan
Belize	El Salvador	Latvia	Palau	Thailand
Benin	Equatorial Guinea	Lesotho	Panama	Timor-Leste
Bhutan	Eritrea	Liberia	Papua New Guinea	Togo
Bolivia (Plurinational State of)	Estonia	Libya	Paraguay	Trinidad and Tobago
Bosnia and Herzegovina	Ethiopia	Lithuania	Peru	Tunisia
Botswana	Fiji	Madagascar	Philippines	Turkmenistan
Brazil	French Polynesia	Malawi	Poland	Tuvalu
Brunei Darussalam	Gabon	Malaysia	Portugal	Uganda
Bulgaria	Gambia	Maldives	Qatar	Ukraine
Burkina Faso	Georgia	Mali	Republic of Korea	United Republic of
Burundi	Ghana	Marshall Islands	Republic of Moldova	Tanzania
Cabo Verde	Greenland	Mauritania	Romania	Uruguay
Cambodia	Guam	Mauritius	Russian Federation	Uzbekistan
Cameroon	Guatemala	Mexico	Rwanda	Vanuatu
Central African Republic	Guinea	Micronesia (Federated States	Saint Vincent and the	Venezuela (Bolivarian
Chad	Guinea-Bissau	of)	Grenadines	Republic of)
China	Guyana	Mongolia	São Tomé and Príncipe	Viet Nam
China, Hong Kong SAR	Haiti	Montenegro	Senegal	Yemen
China, Macao SAR	Honduras	Morocco	Serbia	Zambia
Colombia	India	Mozambique	Seychelles	Zimbabwe
Comoros	Indonesia	Myanmar	Sierra Leone	

Source: World Health Organization Global Health Observatory, Tuberculosis Incidence 2014. Countries with incidence rates of ≥ 20 cases per 100,000 population. For future updates, refer to <http://www.who.int/tb/country/en/>.

Have you had frequent or prolonged visits* to one or more of the countries or territories listed above with a high prevalence of TB disease? (If yes, CHECK the countries or territories, above) Yes No

Have you been a resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)? Yes No

Have you been a volunteer or health care worker who served clients who are at increased risk for active TB disease? Yes No

Have you ever been a member of any of the following groups that may have an increased incidence of latent *M. tuberculosis* infection or active TB disease: medically underserved, low-income, or abusing drugs or alcohol? Yes No

If the answer is YES to any of the above questions, Bryant University requires that you receive TB testing as soon as possible but at least prior to the start of the subsequent semester).

If the answer to all the above questions is NO, no further testing or further action is required.

* The significance of the travel exposure should be discussed with a health care provider and evaluated.

Part II. Clinical Assessment by Health Care Provider

Clinicians should review and verify the information in Part I. Persons answering YES to any of the questions in Part I are candidates for either Mantoux tuberculin skin test (TST) or Interferon Gamma Release Assay (IGRA) unless a previous positive test has been documented.

History of a positive TB skin test or IGRA blood test? (If yes, document below) Yes ____ No ____

History of BCG vaccination? (If yes, consider IGRA if possible.) Yes ____ No ____

1. TB Symptom Check

Does the student have signs or symptoms of active pulmonary tuberculosis disease? Yes ____ No ____

If no, proceed to 2 or 3

If yes, check below:

- Cough (especially if lasting for 3 weeks or longer) with or without sputum production
- Coughing up blood (hemoptysis)
- Chest pain
- Loss of appetite
- Unexplained weight loss
- Night sweats
- Fever

Proceed with additional evaluation to exclude active tuberculosis disease including tuberculin skin testing, chest x-ray, and sputum evaluation as indicated.

2. Tuberculin Skin Test (TST)

(TST result should be recorded as actual millimeters (mm) of induration, transverse diameter; if no induration, write "0". The TST interpretation should be based on mm of induration as well as risk factors.)

Date Given: ___/___/___ Date Read: ___/___/___

Result: _____ mm of induration Interpretation: positive___ negative___

Date Given: ___/___/___ Date Read: ___/___/___

Result: _____ mm of induration Interpretation: positive___ negative___

3. Interferon Gamma Release Assay (IGRA)

Date Obtained: ___/___/___ (specify method) QFT-GIT T-Spot other___

Result: negative___ positive___ indeterminate___ borderline___ (T-Spot only)

4. Chest x-ray: (Required if TST or IGRA is positive)

Date of chest x-ray: ___/___/___ Result: normal___ abnormal___

Part III. Management of Positive TST or IGRA

All students with a positive TST or IGRA with no signs of active disease on chest x-ray should receive a recommendation to be treated for latent TB with appropriate medication. However, students in the following groups are at increased risk of progression from LTBI to TB disease and should be prioritized to begin treatment as soon as possible.

- Infected with HIV
- Recently infected with *M. tuberculosis* (within the past 2 years)
- History of untreated or inadequately treated TB disease, including persons with fibrotic changes on chest radiograph consistent with prior TB disease
- Receiving immunosuppressive therapy such as tumor necrosis factor-alpha (TNF) antagonists, systemic corticosteroids equivalent to/greater than 15 mg of prednisone per day, or immunosuppressive drug therapy following organ transplantation
- Diagnosed with silicosis, diabetes mellitus, chronic renal failure, leukemia, or cancer of the head, neck, or lung
- Have had a gastrectomy or jejunioileal bypass
- Weigh less than 90% of their ideal body weight
- Cigarette smokers and persons who abuse drugs and/or alcohol

••Populations defined locally as having an increased incidence of disease due to *M. tuberculosis*, including medically underserved, low-income populations

_____ Student agrees to receive treatment

_____ Student declines treatment at this time

Health Care Professional Signature

Date